

a. distance between rows and/or plants :

1. Variety name :	AGERATUM QIS BOUQUET WHITE
2. Description : Our own selection of open pollinated Age Bouquet White is branching when pinche with a tuft of white petals.	eratum for fresh and dried cutflower use. ed and carries a cluster of button-formed flowers
3. General a. colors b. height c. life cycle d. weight of 1000 seeds e. time sowing/flowering f. family g. hardiness zone	clear white 60 cm annual 0,2 gr; 0,35 gr for 1000 plants 16-20 weeks Compositae
4. Sowing instructions a. method b. time c. temperature d. germination time e. seed quantity or plants per m ²	transplant plant habit is practically daylength neutral. Therefore Ageratum is cultivated in greenhouses throughout the year. In the open Ageratum can be transplanted from end of April until mid June. Time from sowing to transplanting is 26 to 35 days, depending on sowing period. 20-22°C 7-14 days when pinched in summer 32 plants per m²; in autumn
5. Growing on	24 plants per m ² . Also non-pinched cultures of 64 plants per m ² are commonly practiced.

32 pl/m² = 18x18 24 pl/m² = 20x20 64 pl/m² = 12,5x12,5

12,5x12,5 cm.

Always produced in wire netting! Preferably

code: 0160.1600

b. soil : any well-drained soil with a normal fertilization level

can be used

c. irrigation : until the flowerheads appear overhead irrigation can

be used, afterwards soil irrigation or flooding should

be used

d. optimum temperature : 15-18°C

e. diseases/pests : to avoid Pythium after transplanting, the soil should

be treated with a preventive fungicide

f. fertilization : a maintainance fertilization of 20 gr of 12-10-18 per

m² of an N-P-K fertilizer is used. To strengthen plants, an extra gift of potassium is advised. Please always remember to water abundantly after any

fertilizer gift over the growing plants.

6. General advice

a. cutting period : 4-5 weeks

b. cutting stage : when flowers are nearly full flowering

c. post delivery treatment : once cut, the flowers should remain at least 1 night

on water before transport or sale. The flowers will then open better and will be stronger. No special

transport treatment is necessary.

d. general : when early sown/planted outdoors two harvest

periods are possible if Ageratum is uniformly cut back

after the first harvest.

This information is based on West European conditions and is given for general guidance only. No guarantee is given for the result of the crop, nor is liability accepted for the consequences of applying above indications.