

syngenta flowers

Dianthus Super Parfait™

Culture Guide

Botanical name: Dianthus chinensis

Product form: Seed

Containers: Packs, Pints, Quarts, Gallons, Hanging

Baskets

Habit: Mounded

Garden Specifications

Garden Height: 6–8" (15–20 cm) tall **Garden Width:** 8–10" (20–25 cm) wide

Exposure: Full sun **USDA zone:** 5 **AHS zone:** 9–1

Product use: Packs, Containers, Hanging Baskets,

Landscapes, Combos Germination

Stages 1 & 2

Germination time: 5 days **Media temp:** 72–74 °F (22–23 °C)

Chamber: Optional

Light: Not required for germination

Seed cover: Yes

Moisture level: 4 - WET: Soil is dark brown but not shiny, no free water is seen at the surface of the soil, when pressed or squeezed water drips easily, and trays are heavy with a visible bend in the middle.

Rec. tray size: 288-cell tray

Seeds per cell: 1

Young Plant Production

Stages 3 & 4 **Temperature**

Day: 66–68 °F (19–20 °C) **Night:** 66–68 °F (19–20 °C)

Average daily temperature: 67 °F (19 °C)

Lighting

Day extension lighting: Beneficial to 14 hours **Light intensity:** 3,500–4,500 foot candles (700–900

Day length response: Facultative long day **Daily light integral:** 12–14 mols/day

Media pH: 5.5-5.9

Media EC: SME EC: 0.9-1.3 mS/cm

Fertilizer: 75-125 ppm N

Moisture level: Media should be allowed to dry between irrigations. Alternate between moisture level 3 and 4.

3 - MOIST: Soil is brown in color, strongly squeezing the soil will extract a few drops of water, and trays are light with no visible bend.

4 - WET: Soil is dark brown but not shiny, no free water is seen at the surface of the soil, when pressed or squeezed water drips easily, and trays are heavy with a visible bend in the middle.

Pinching: No

Plant growth regulators (PGRs): If needed, spray with Bonzi[®] at 2–5 ppm or Sumagic[®] at 3–5 ppm.

Plug grow time: 5–6 weeks in a 288-cell tray **Comments:** Providing high light and dropping the temperature to an ADT of 60 °F (21 °C) will reduce the need for PGRs. Day extension lighting to 14 hours will hasten flowering of the finished plant.

Finishing

Temperature

Day: 62–64 °F (17–18 °C) **Night:** 62–64 °F (17–18 °C)

Average daily temperature: 63 °F (17 °C)

Lighting

Day extension lighting: Beneficial to 14 hours **Light intensity:** 3,500–4,500 foot candles (700–900

micro mols)

Day length response: Facultative long day **Daily light integral:** 12–14 mols/day

Transplanting: Transplant directly into the finished container. Place the rooting media slightly above the level of media in the container. Make sure that the root ball is covered and that the plug is situated in the center of the pot.

Media pH: 5.5-5.9

Media EC: SME EC: 1.5–2.1 mS/cm, PourThru EC: 2.3–3.2

mS/cm

Fertilizer: 125-175 ppm N

Pinching: No. Pinching is not recommended or needed.

Plant growth regulators (PGRs): If needed, spray with Bonzi® at 5-8 ppm.

Tech tip: Providing high light and growing cool will reduce the need for PGRs. Dianthus also responds well to a 5–10 °F (3–6 °C) negative DIF in the early morning. Day extension lighting to 14 hours will hasten flowering.



CHRYSAL Try Chrysal Alesco®, a postharvest foliar spray, to protect ethylene sensitive crops during shipping and retail

Moisture level: Media should be allowed to dry between irrigations. Alternate between moisture level 3 and 4.

- 3 MOIST: Soil is brown in color, strongly squeezing the soil will extract a few drops of water, and trays are light with no visible bend.
- 4 WET: Soil is dark brown but not shiny, no free water is seen at the surface of the soil, when pressed or squeezed water drips easily, and trays are heavy with a visible bend in the middle.

Common pests: Aphids, Thrips, Spider mites **Common diseases:** Fusarium, Alternaria, Botrytis

Scheduling

Size	Crop Time	Plants Per Pot
Packs	9-11 weeks	1 ррр
Large Packs	9-11 weeks	1 ррр
1.0 pint (4 inch)	9-11 weeks	1 ppp
1.0 quart (4.5 to 5 inch)	10-12 weeks	1–2 ppp
1.25 to 2.5 quart (5.5 to 6.5 inch, trade gallon)	10-12 weeks	3 ррр
1.5 gallon hanging basket (10 inch basket)	10-12 weeks	4–5 ppp

Estimated finish crop time is from transplant of a 288-cell tray and finished at an average daily temperature (ADT) of 63 °F (17 °C).

Example crop schedule for a large pack

Weeks From Transplant	Description
1 week	Provide day extension lighting to 14 hours to hasten flowering
4 weeks	If needed spray with Bonzi® at 5–8 ppm
7 weeks	If needed spray with Bonzi® at 5–8 ppm
10 weeks	Ready to ship



syngenta.flowers

All photos are either the property of Syngenta or are used with permission.

© 2020 Syngenta. Important: Always read and follow label instructions. Some products may not be registered for sale or use in all states or counties. Please check with your local Extension Service to ensure registration status. Some or all of the varieties may be protected under one or more of the following: Plant Variety Protection, United States Plant Patents and/or Utility Patents and may not be propagated or reproduced without authorization. The trademarks or service marks displayed or otherwise used herein are the property of a Syngenta Group Company. All other trademarks are the property of their respective owners